MycoKeys 86: 47–63 (2022)
doi: 10.3897/mycokeys.86.71623
https://mycokeys.pensoft.net

RESEARCH ARTICLE

A peer-reviewed open-access journal
MycoKeys
Launched to accelerate biodiversity research

Three new *Xylaria* species (Xylariaceae, Xylariales) on fallen leaves from Hainan Tropical Rainforest National Park

Xiao-Yan Pan[1,2], Zi-Kun Song[2,3], Zhi Qu[2], Tie-Dong Liu[1], Hai-Xia Ma[2,4]

1 *College of Forestry, Hainan University, Haikou 570228, China* **2** *Institute of Tropical Bioscience and Biotechnology, Chinese Academy of Tropical Agricultural Sciences, Haikou 571101, China* **3** *College of Plant Protection, Jilin Agricultural University, Jilin 130000, China* **4** *Hainan Institute for Tropical Agricultural Resources, Chinese Academy of Tropical Agricultural Sciences, Haikou 571101, China*

Corresponding authors: Hai-Xia Ma (mahaixia@itbb.org.cn), Tie-Dong Liu (liu@hainanu.edu.cn)

Academic editor: Thorsten Lumbsch | Received 15 July 2021 | Accepted 21 December 2021 | Published 12 January 2022

Citation: Pan X-Y, Song Z-K, Qu Z, Liu T-D, Ma H-X (2022) Three new *Xylaria* species (Xylariaceae, Xylariales) on fallen leaves from Hainan Tropical Rainforest National Park. MycoKeys 86: 47–63. https://doi.org/10.3897/mycokeys.86.71623

Abstract

Three new species of *Xylaria* on fallen leaves in Hainan Province of China are described and illustrated, based on morphological and molecular evidence. *Xylaria hedyosmicola* is found on fallen leaves of *Hedyosmum orientale* and featured by thread-like stromata with a long sterile filiform apex. Phylogenetically, *X. hedyosmicola* is closely related to an undescribed *Xylaria* sp. from Hawaii Island, USA and morphologically similar to *X. vagans*. *Xylaria lindericola* is found on fallen leaves of *Lindera robusta* and characterised by its subglobose stromata and a long filiform stipe. It is phylogenetically closely related to *X. sicula* f. *major*. *Xylaria polysporicola* is found on fallen leaves of *Polyspora hainanensis*, it is distinguished by upright or prostrate stromata and ascospores sometimes with a slimy sheath or non-cellular appendages. *Xylaria polysporicola* is phylogenetically closely related to *X. amphithele* and *X. ficicola*. An identification key to the ten species on fallen leaves in China is given.

Keywords

Folicolous fungi, Phylogeny, Pyrenomycetes, Taxonomy

Copyright *Xiao-Yan Pan et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

Species of *Xylaria* Hill ex Schrank are commonly found throughout the temperate, subtropical and tropical regions of the world, associated with wood, fallen fruits or seeds, fallen leaves or petioles and termite nests (Dennis 1956; Rogers 1986; Rogers and Samuels 1986; San Martin and Rogers 1989; Ju and Rogers 1999; Ju and Hsieh 2007; Fournier 2014). Previous studies on *Xylaria* have dealt primarily with species growing on wood and termite nests (Rogers et al. 2005; Ju and Hsieh 2007; Fournier et al. 2020), but the species diversity and distribution of the genus on other substrates, such as fallen fruits or seeds and fallen leaves or petioles, are still poorly studied (Hsieh et al. 2010; Ju et al. 2018). Especially, the study of *Xylaria* species growing on fallen leaves or petioles is far behind those mentioned taxa associated with other substrates and only seven species have been reported on those substrates in China (Dennis 1956; Rogers et al. 1988; Zhu and Guo 2011; Huang et al. 2014, 2015; Ma and Li 2018).

Hainan Province (20°01.04'N, 110°20.95'E) is located in southern China and enjoys a tropical monsoon climate. More than 6036 plant species, 1895 genera and 243 families have been reported in the province (Yang 2015). Different kinds of tropical vegetations (e.g. Moraceae, Euphorbiaceae and Arecaceae) and rainforests are distributed over the vast territory of the province, in which abundant fungi occur (Dai et al. 2009; Dai 2012; Gao and Yang 2016; Cui et al. 2019). Two intensive surveys of xylariaceous fungi were carried out in Hainan province in 2019 and 2020 and about 400 specimens of Xylariaceae were collected. These materials have been carefully studied through both morphological and phylogenetic methods and three new species on fallen leaves were identified. The new taxa are described and illustrated, and an identification key is provided for the 10 known species of *Xylaria* on fallen leaves in China.

Materials and methods

Morphological studies

Voucher specimens are deposited in the Fungarium of the Institute of Tropical Bioscience and Biotechnology, Chinese Academy of Tropical Agricultural Sciences (FCATAS), Hainan Province, China. Samples for microscopic examination were mounted in distilled water, Melzer's reagent, India ink or 1% SDS. Microscopic features observation, measurements and photographing were performed by using a Zeiss Axio Imager A2 microscope (Göttingen, Germany) by differential interference contrast microscopy (DIG) and brightfield microscopy (BF). The photographs of stromata, perithecia and ostioles were taken with a VHX-600E stereomicroscope Keyence Corporation (Osaka Japan). The methods of collecting, preservation and identification of the specimens follow Ma and Li (2018).

DNA extraction and sequencing

A modified cetyltrimethylammonium bromide (CTAB) extraction kit (Aidlab Biotechnologies, Beijing, China) was employed for total DNA extraction from dried specimens. The ITS region was amplified with the primer pair ITS4 and ITS5 (White et al. 1990) using the following procedure: initial denaturation at 95 °C for 3 min, followed by 30 cycles of 94 °C for 40 s, 55.8 °C for 45 s and 72 °C for 1 min and a final extension of 72 °C for 10 min. The TUB and RPB2 gene region were amplified with primers T1/T22 (O'Donnell and Cigelnik 1997) and fRPB2-5F/fRPB2-7CR (Liu et al. 1999), respectively, using the following procedure: initial denaturation at 95 °C for 3 min, followed by 35 °C cycles of 94 °C for 1 min, 52 °C for 1 min and 72 °C for 1.5 min and a final extension of 72 °C for 10 min (Hsieh et al. 2005). DNA sequencing was performed at BGI tech (Guangzhou, China) and sequences were deposited in GenBank (Table 1).

Table 1. Species, specimens and GenBank accession number of sequences used in this study. New species and sequences are set in bold.

Taxon	Substrate / Origin	Specimen No.	GenBank No.			Reference
			ITS	TUB	RPB2	
Xylaria acuminatilongissima	termite nests / China Taiwan	HAST 623	EU178738	GQ502711	GQ853028	Hsieh et al. (2010)
X. adscendens	wood / Guadeloupe	HAST 570	GU300101	GQ487708	GQ844817	Hsieh et al. (2010)
*X. allantoide*a	trunk / China Taiwan	HAST 94042903	GU324743	GQ502692	GQ848356	Hsieh et al. (2010)
X. amphithele	dead leaves / Guadeloupe	HAST 529	GU300083	GQ478218	GQ844796	Hsieh et al. (2010)
X. apoda	bark / China Taiwan	HAST 90080804	GU322437	GQ495930	GQ844823	Hsieh et al. (2010)
X. arbuscula	bark / China Taiwan	HAST 89041211	GU300090	GQ478226	GQ844805	Hsieh et al. (2010)
X. arbuscula var. *plenofissura*.	wood / China Taiwan	HAST 93082814	GU339495	GQ478225	GQ844804	Hsieh et al. (2010)
X. atrodivaricata	termite nests / China Taiwan	HAST 95052001	EU178739	GQ502713	GQ853030	Hsieh et al. (2010)
X. badia	bamboo culm / China Taiwan	HAST 95070101	GU322446	GQ495939	GQ844833	Hsieh et al. (2010)
X. bambusicola	bamboo culm / Thailand	JDR 162	GU300088	GQ478223	GQ844801	Hsieh et al. (2010)
X. berteri	bark / USA	JDR 256	GU324750	GQ502698	GQ848363	Hsieh et al. (2010)
X. berteri	bark / China Taiwan	HAST 90112623	GU324749	AY951763	GQ848362	Hsieh et al. (2010)
X. betulicola	leaves of *Betula* / China	FCATAS 750	MF774332	–	–	Ma and Li (2018)
X. brunneovinosa	termite nests / China Taiwan	HAST 720	EU179862	GQ502706	GQ853023	Hsieh et al. (2010)
X. castorea	wood / New Zealand	PDD 600	GU324751	GQ502703	GQ853018	Hsieh et al. (2010)
X. cirrata	termite nests / China Taiwan	HAST 664	EU179863	GQ502707	GQ853024	Hsieh et al. (2010)
X. coccophora	wood / French	HAST 786	GU300093	GQ487701	GQ844809	Hsieh et al. (2010)
X. crinalis	wood / China	FCATAS 751	MF774330	–	–	Ma and Li (2018)
X. crozonensis	bark / France	HAST 398	GU324748	GQ502697	GQ848361	Hsieh et al. (2010)
X. cubensis	log / Russian Far East	HAST 477	–	GQ502699	GQ848364	Hsieh et al. (2010)
X. culleniae	pod / Thailand	JDR 189	GU322442	GQ495935	GQ844829	Hsieh et al. (2010)
X. escharoidea	termite nests / China Taiwan	HAST 658	EU179864	GQ502709	GQ853026	Hsieh et al. (2010)
X. feejeensis	bark / China Taiwan	HAST 92092013	GU322454	GQ495947	GQ848336	Hsieh et al. (2010)
X. ficicola	fallen leaves and petioles of *Ficus auriculata* / China	HMJAU 22818	MZ351258	–	–	**This study**
X. filiformis	herbaceous stem / Iran	GUM 1052	KP218907	–	–	Hashemi et al. (2015)
X. fimbriata	termite nests / French West Indies	HAST 491	GU324753	GQ502705	GQ853022	Hsieh et al. (2010)
X. cf. *glebulosa*	fruit / French West Indies	HAST 431	GU322462	GQ495956	GQ848345	Hsieh et al. (2010)

Taxon	Substrate / Origin	Specimen No.	GenBank No.			Reference
			ITS	TUB	RPB2	
X. grammica	wood / China Taiwan	HAST 479	GU300097	GQ487704	GQ844813	Hsieh et al. (2010)
X. griseosepiacea	termite nests / China Taiwan	HAST 641	EU179865	GQ502714	GQ853031	Hsieh et al. (2010)
X. hedyosmicola	**fallen leaves of *Hedyosmum orientale* / China Hainan**	**FCATAS 856 (HT)**	**MZ227121**	**MZ221183**	**MZ683407**	**This study**
X. hedyosmicola	**fallen leaves of *Hedyosmum orientale* / China Hainan**	**FCATAS 857**	**MZ227023**	**MZ221184**	**MZ851780**	**This study**
X. hypoxylon	wood / Belgium	HAST 152	GU300096	GQ260187	GQ844812	Hsieh et al. (2010)
X. hypoxylon	wood / China Taiwan	HAST 95082001	GU300095	GQ487703	GQ844811	Hsieh et al. (2010)
X. hypoxylon	leaf debris / Sweden	CBS 122617	AM993146	–	–	Persoh et al. (2009)
X. ianthinovelutina	fruit of *Swietenia* / Martinique	HAST 553	GU322441	GQ495934	GQ844828	Hsieh et al. (2010)
X. intraflava	termite nests / China Taiwan	HAST 725	EU179866	GQ502718	GQ853035	Hsieh et al. (2010)
X. juruensis	*Arenga engleri* / China Taiwan	HAST 92042501	GU322439	GQ495932	GQ844825	Hsieh et al. (2010)
X. laevis	wood / Martinique	HAST 419	GU324746	GQ502695	GQ848359	Hsieh et al. (2010)
X. leavis	bark / China Taiwan	HAST 95072910	GU324747	GQ502696	GQ848360	Hsieh et al. (2010)
X. lindericola	**fallen leaves of *Lindera robusta* / China Hainan**	**FCATAS 852 (HT)**	**MZ005635**	**MZ031978**	**MZ031982**	**This study**
X. lindericola	**fallen leaves of *Lindera robusta* / China Hainan**	**FCATAS 853**	**MZ005636**	**MZ031979**	**MZ048749**	**This study**
X. liquidambar	fruits of *Liquidambar formosana* / China Taiwan	HAST 93090701	GU300094	GQ487702	GQ844810	Hsieh et al. (2010)
X. longissima	wood / China	FCATAS 749	MF774331	–	–	Ma and Li (2018)
X. longissima	wood / Iran	IRAN 16582 F	KP218906	–	–	Hashemi et al. (2015)
X. meliacearum	petioles and infructescence of *Guarea guidonia* / Puerto Rico	JDR 148	GU300084	GQ478219	GQ844797	Hsieh et al. (2010)
X. multiplex	wood / USA	JDR 259	GU300099	GQ487706	GQ844815	Hsieh et al. (2010)
X. muscula	dead branch / French West	HAST 520	GU300087	GQ478222	GQ844800	Hsieh et al. (2010)
X. nigripes	termite nests / China Taiwan	HAST 653	GU324755	GQ502710	GQ853027	Hsieh et al. (2010)
X. oxyacanthae	fallen seeds / USA	JDR 859	GU322434	GQ495927	GQ844820	Hsieh et al. (2010)
X. oxyacanthae	fruits / Germany	LZ 2010-502	HQ414587	–	–	Roensch et al. (2010)
X. palmicola	fruits / New Zealand	PDD 604	GU322436	GQ495929	GQ844822	Hsieh et al. (2010)
X. phyllocharis	dead leaves / French West	HAST 528	GU322445	GQ495938	GQ844832	Hsieh et al. (2010)
X. plebeja	trunk / China Taiwan	HAST 91122401	GU324740	GQ502689	GQ848353	Hsieh et al. (2010)
X. polymorpha	wood / USA	JDR 1012	GU322460	GQ495954	GQ848343	Hsieh et al. (2010)
X. polymorpha	Stump / Germany	M:M-0125909	FM164944	–	–	Persoh et al. (2009)
X. polysporicola	**fallen leaves of *Polyspora hainanensis* / China Hainan**	**FCATAS 848 (HT)**	**MZ005592**	**MZ031976**	**MZ031980**	**This study**
X. polysporicola	**fallen leaves of *Polyspora hainanensis* / China Hainan**	**FCATAS 849**	**MZ005591**	**MZ031977**	**MZ031981**	**This study**
X. regalis	log of *Ficus racemose* / India	HAST 920	GU324745	GQ502694	GQ848358	Hsieh et al. (2010)
X. schweinitzii	bark / China Taiwan	HAST 92092023	GU322463	GQ495957	GQ848346	Hsieh et al. (2010)
X. sicula f. *major*	fallen leaves / China Taiwan	HAST 90071613	GU300081	GQ478216	GQ844794	Hsieh et al. (2010)
Xylaria sp. 6	fallen leaves of *Tibouchina semidecandra* / USA	JDR 258	GU300082	GQ478217	GQ844795	Hsieh et al. (2010)
X. striata	branch / China	HAST 304	GU300089	GQ478224	GQ844803	Hsieh et al. (2010)
X. tentaculata	leaf litter or wood / Korea	KA12-0530	KM077162	–	–	Kim et al. (2016)
X. tentaculata	leaf litter or wood / Korea	KA13-1324	KM077163	–	–	Kim et al. (2016)
X. tentaculata.	leaf litter or wood / Korea	KA13-1325	KM077164	–	–	Kim et al. (2016)
X. venosula	twigs / USA	HAST 94080508	EF026149	EF025617	GQ844806	Hsieh et al. (2010)
X. venustula	bark / China Taiwan	HAST 88113002	GU300091	GQ487699	GQ844807	Hsieh et al. (2010)
X. xylarioides	wood / Iran	GUM 1151	KP218909	–	–	Hashemi et al. (2015)
Hypoxylon fragiforme	bark / France	HAST 383	JN979420	AY951720	–	Hsieh et al. (2005)
Camillea obularia	– / Puerto Rico	ATCC 28093	KY610384	KX271243	–	Wendt et al. (2018)

Phylogenetic analyses

The molecular phylogeny was inferred from a combined dataset of ITS, TUB and RPB2 sequences. The sequences retrieved from open databases originated from Hsieh et al. (2005), Persoh et al. (2009), Hsieh et al. (2010), Roensch et al. (2010), Hashemi et al. (2015), Kim et al. (2016), Ma and Li (2018) and Wendt et al. (2018) (Table 1). *Hypoxylon fragiforme* (Pers.) J. Kickx f. and *Camillea obularia* (Fr.) Læssøe, J.D. Rogers & Lodge were selected as outgroup taxa. Sequences were aligned using the MAFFT online (http://mafft.cbrc.jp/alignment/server/). Alignments were optimised manually in BioEdit 7.0.5.3 (Hall 1999).

A combined matrix of ITS-RPB2-TUB and ITS-exons of TUB and RPB2 were used to construct phylogenetic analysis by two methods including maximum likelihood (ML) and Bayesian Inference (BI) analysis, respectively. ML tree generation and bootstrap analyses were performed via the programme RAxML7.2.6 (Stamatakis 2006) running 1000 replicates combined with a ML search. Bayesian analysis was performed with MrBayes 3.1 (Huelsenbeck and Ronquist 2005) implementing the Markov Chain Monte Carlo (MCMC) technique and parameters predetermined by MrModeltest 2.3 (Nylander 2004).

Results

Molecular phylogeny

This study used genetic sequences of 57 species, including 69 ITS sequences, 57 TUB sequences and 54 RPB2 sequences. We applied two tree construction methods to improve the reliability of the results.

After the alignment sequence was adjusted using MAFFT, the ITS alignment, shown in BioEdit 7.0.5, consisted of 778 character positions, 2219 in the TUB alignment and 1241 in the RPB2 alignment. After curing, the constructed multigene alignment (MGA) consisted of 3138 characters (523 of which were derived from the ITS alignment, 1550 from TUB alignment, 1065 from RPB2 alignment). Of the MGA, 1354 characters were considered parsimony-informative.

The analysis results show that the phylogenetic tree, generated by ML in RAxML7.2.6, is basically the same as that generated by BI in MrBayes 3.1. Topology of the phylogenetic analyses, based on ITS-RPB2-TUB and ITS-exons of TUB and RPB2, have no significant conflicts. Only the BI tree is shown in Figure 1 with Bayesian posterior probabilities ≥ 0.95 and ML bootstrap values ≥ 50% labelled along the branches. The phylogenetic tree showed that *X. hedyosmicola* is clustered with *Xylaria* sp. 6, *X. polysporicola* is clustered with *X. amphithele* F. San Martín & J.D. Rogers and *X. ficicola* Hai X. Ma, Lar.N. Vassiljeva & Yu Li, *X. lindericola* is clustered with *X. sicula* Pass. & Beltr. f. *major* Ciccarone, but were separated from other species, as well as from each other.

Taxonomy

Xylaria hedyosmicola **Hai X. Ma & X.Y. Pan, sp. nov.**
MycoBank No: 839780
GenBank No: MZ227121, MZ221183, MZ683407
Figure 2

Diagnosis. Differs from *X. vagans* by its stromata without a black rhizomorphoid mycelium connecting dead leaves, larger ascospores and tubular to slightly urn-shaped apical apparatus. Differs from *X. betulicola* by its smaller stromta and larger ascospores.

Typification. China. Hainan Province, Lingshui County, Diaoluoshan Natural Reserve, on fallen leaves of *Hedyosmum orientale* (Chloranthaceae), 31 December 2020, Haixia Ma (holotype, FCATAS 856).

Etymology. "***hedyosmicola***" refers to the growth on leaves of *Hedyosmum orientale*.

Teleomorph. *Stromata* upright, solitary to cespitose, thread-like, unbranched or occasionally branched once at top, 2–5.5 cm total length; with a long sterile filiform apex up to 0.5–3 cm long; fertile part 3–17 mm long × 0.5–1 mm diam., usually consisting of closely packed or scattered perithecia; stipe 8–18 mm long × 0.1–0.5 mm diam., glabrous, finely longitudinally striate, the base slightly swollen; surface roughened, with half-exposed to fully exposed perithecial contours and wrinkles. Externally black, interior white. Texture soft. *Perithecia* subglobose, 200–470 μm diam. *Ostioles* papillate, 11–22 μm diam. *Asci* with eight ascospores arranged in uniseriate manner, cylindrical, 105–160 μm total length, the spore-bearing parts 70–100 μm long × 8–12 μm broad, the stipes 25–70 μm long, with apical apparatus bluing in Melzer's reagent, tubular to slightly urn-shaped, 2.5–4.8 μm high × 2.5–3.5 μm broad. *Ascospores* brown, unicellular, ellipsoid-inequilateral, with narrowly rounded ends, smooth, (12–)13–15(–16.7) × (6–) 6.5–7.5 (–8.5) μm (M = 14 × 7 μm, n = 60), straight to slightly sigmoid germ slit spore-length or almost spore-length, with a slimy sheath on ventral side swollen at both ends to form rounded non-cellular appendages visible in Indian ink.

Additional specimen examined. China. Hainan Province, Lingshui County, Diaoluoshan Natural Reserve, on fallen leaves of *Hedyosmum orientale*, 31 December 2020, Haixia Ma (FCATAS 857).

Remarks. *Xylaria hedyosmicola* closely resembles *X. vagans* Petch by sharing thread-like or long hair-like stromata bearing closely packed or scattered perithecia with a long sterile filiform apex. *Xylaria vagans* was originally described and illustrated by Petch (1915) from Sri Lanka. However, based on comparisons of the descriptions and illustrations, there were some differences between the two species. *Xylaria hedyosmicola* has larger sporiferous part of asci (70–100 μm × 8–12 μm) with tubular to slightly urn-shaped apical apparatus bluing in Melzer's reagent, brown and larger ascospores with straight (Fig. 2n and p) to slightly sigmoid germ slit (Fig. 2o), with narrowly rounded ends and a slimy sheath on ventral side swollen at both ends to form rounded non-cellular appendages, while *X. vagans* has a black rhizomorphoid mycelium connecting dead leaves, smaller sporiferous part 68–72 μm × 6 μm and black-brown, cymbiform,



Figure 1. Phylogenetic tree of *Xylaria* based on multigene alignment of ITS-TUB-RPB2 in the Bayesian analysis. Bayesian posterior probabilities (≥ 0.95, before the slash markers) and RaxML bootstrap values (≥ 50, after the slash markers) are shown. Different clades are indicated as coloured blocks.

smaller ascospores 9–12 × 5–6 µm, with broadly rounded ends and is without apical apparatus, germ slit and sheath or appendages (Petch 1915). Unfortunately, the molecular sequences of *X. vagans* from Sri Lanka were not available.

Xylaria betulicola Hai X. Ma, Lar.N. Vassiljeva & Yu Li is similar to *X. hedyosmicola* in stromatal morphology, but differs in having larger stromata 3–7 cm, slightly smaller



Figure 2. *Xylaria hedyosmicola* (FCATAS 856, holotype) **a, b, e** stromata on leaves (b, FCATAS 857) **c** stromatal surface **d** section through stroma, showing a perithecium **f** immature asci in water **g, h** ascal apical ring in Melzer's reagent **i, j** ascospores in Melzer's reagent **k** ascus in 1% SDS **l, m** asci and ascal apical ring in Melzer's reagent **n** ascospore in Melzer's reagent showing straight germ slit **o** ascospore in Melzer's reagent showing slightly sigmoid germ slit **p, q** ascospore showing a slimy sheath and non-cellular appendages in India ink. Scale bars: 1 cm (**a, b**); 0.1 mm (**c, d**); 0.5 mm (**e**); 20 μm (**f, m**); 10 μm (**g–l, n–q**).

ascospores (11.5)12–14(15) × 5–6 μm, without sheath or appendages (Ma and Li 2018). In the phylogenetic tree, *X. hedyosmicola* formed a fully supported clade with *Xylaria* sp. 6 from Hawaiian Islands, USA (Hsieh et al. 2010). Although there are no descriptions on *Xylaria* sp. 6 in the study of Hsieh et al. (2010), we suspected that it is conspecific with *X. hedyosmicola*. The sequences comparison showed that there are 98.7%, 99% and 99.9% maximal percentage identities, respectively in ITS, TUB and RPB2 between *X. hedyosmicola* (FCATAS 856) and *Xylaria* sp. 6 from USA (JDR 258).

Xylaria lindericola **Hai X. Ma & X.Y. Pan, sp. nov.**
MycoBank No: 839554
GenBank No: MZ005635, MZ031978, MZ031982
Figure 3

Diagnosis. Differs from *X. sicula* f. *major* by its subglobose stromata without a long sterile apex, larger ascospores and host plant. Differs from *X. hypsipoda* by its black stromata, glabrous stipes and smaller apical apparatus.

Typification. China. Hainan Province, Lingshui County, Diaoluoshan Natural Reserve, on fallen leaves of *Lindera robusta* (Lauraceae), 31 December 2020, Haixia Ma (holotype, FCATAS 852).

Etymology. "***lindericola***" refers to the growth on leaves of *Lindera robusta*.

Teleomorph. *Stromata* upright or prostrate, solitary to cespitose, unbranched or branched once or more at stipe, 3–26 cm total length; fertile part subglobose on long filiform stipes, 0.1–0.4 cm diam., the stipe 3–25 cm long × 0.1–1 mm diam., glabrous, finely longitudinally striate, the base slightly swollen; surface roughened by wrinkles and barely exposes perithecial contours. External black, interior white. Texture soft. *Perithecia* subglobose, 300–550 μm diam. *Ostioles* black, papillate. *Asci* with eight ascospores in uniseriate manner, cylindrical, 105–165 μm total length, the spore-bearing parts 65–115 μm long × 7.5–10.5 μm broad, the stipes 25–65 μm long, with apical apparatus bluing in Melzer's reagent, tubular to urn-shaped, 3.9–5.5 μm high × 3–5 μm broad. *Ascospores* brown, unicellular, ellipsoid-inequilateral, with slightly narrowly rounded ends, aberrant ascospores with strongly pinched or beaked ends, smooth, (12.5–)13.5–15.5(–18) × (7–) 7.5–8.5 (–9.5) μm (M = 14.8 × 8 μm, n=60), with straight germ slit spore-length, without sheath or appendages visible in India ink.

Additional specimen examined. China. Hainan Province, Lingshui County, Diaoluoshan Natural Reserve, on fallen leaves of *Lindera robusta*, 31 December 2020, Haixia Ma (FCATAS 853).

Remarks. *Xylaria lindericola* is distinguished by its subglobose fertile part of stroma on a long filiform stipe and growing on fallen leaves of *Lindera robusta*. The species is somewhat similar to *X. sicula* f. *major* in morphology of stromatal fertile part. However, *X. sicula* f. *major* has stromata with long sterile apex, slightly smaller ascospores 9–13(–15) × (3–) 4.5–6 (–7) μm and grows on dead *Olea* leaves (Ciccarone 1947;



Figure 3. *Xylaria lindericola* (FCATAS 852, holotype) **a, b** stromata on leaves **c** fertile part of stroma **d** stromatal surface **e** section through stroma, showing perithecia **f** ascal apical ring and ascospores with beaked ends in Melzer's reagent **g** ascus and ascal apical ring in Melzer's reagent **h** ascus in water **i, j** ascospores in water **k, l** ascospore in Melzer's reagent **m** ascospore in India ink **n** ascospore in 1% SDS showing germ slit. Scale bars: 1.5 cm (**a, b**); 0.2 mm (**c–e**); 10 μm (**f–n**).

Graniti 1959; Fournier 2014). In the phylogenetic tree, *X. lindericola* formed a fully supported clade with *X. sicula* f. *major* (Figure 1).

Xylaria hypsipoda Massee is similar to *X. lindericola* by sharing globose stromata and ascospores dimensions, but differs in having stromata with whitish scales, hairy stipes and urn-shaped, slightly larger apical apparatus 5–8 μm high × 2.9–5 μm broad (Rogers et al. 1987).

Xylaria ficicola resembles *X. lindericola* in stromatal morphology, but differs in having strongly exposed perithecial mounds of stromatal surface, larger ascospores (16–) 17.5–21(–22.7) × 6.5–8.5 μm with conspicuous hyaline noncellular appendage and grows on fallen leaves and petioles of *Ficus auriculata* (Ma et al. 2011). *Xylaria heloidea* Penz. & Sacc. from Indonesia is somewhat similar to *X. lindericola* in stromatal morphology, but the former has obconical, convex stromatal top, larger ascospores (14.5–) 15.5–18(–19) × (5–)5.5–6.5(–7) μm (16.7 × 6.1 μm), with a hyaline sheath swelling at both ends to form non-cellular appendages and grows on fallen fruits, twigs, petioles, and leaves of various plants (Ju et al. 2018).

Xylaria comosa (Mont.) Fr. and *X. clusiae* K.F. Rodrigues, J.D. Rogers & Samuels are also somewhat similar to *X. lindericola* in stromatal morphology. However, *X. comosa* has larger ascospores (21)–26–40 × 7–11 μm and larger apical ring 10.5 μm high × 7.5 μm broad (Dennis 1956) and *X. clusiae* has smaller stromata 1–3.5 cm, ascospores broadly ovoida1 to nearly globose (11.6–)12.8–16.7(–18) × 8–15 μm, with colorless appendage at one end (Samuels and Rogerson 1990).

Xylaria polysporicola **Hai X. Ma & X.Y. Pan, sp. nov.**
MycoBank No: 839552
GenBank No: MZ005592, MZ031976, MZ031980
Figure 4

Diagnosis. Differs from *X. phyllocharis* by its half-exposed to fully exposed perithecial contours, the fertile part cylindrical and larger perithecia. Differs from *X. phyllophila* by its smaller ascospores. Differs from *X. amphithele* by its cylindrical stromata.

Typification. China. Hainan Province, Lingshui County, Diaoluoshan Natural Reserve, on fallen leaves of *Polyspora hainanensis* (Theaceae), 31 December 2020, Haixia Ma (holotype, FCATAS 848).

Etymology. "***polysporicola***" refers to the growth on leaves of *Polyspora hainanensis*.

Teleomorph. *Stromata* solitary, upright or prostrate, cylindrical, unbranched or occasionally branched, 1–4 cm total length, with acute sterile apex up to 2 mm long; fertile part 2–15 mm long × 0.5–1.6 mm diam., usually consists of closely packed perithecia and occasionally with scattered perithecia; the stipe 5–30 mm long × 0.3–1 mm diam., glabrous, finely longitudinally striate, the base slightly swollen; surface roughened, with half-exposed to fully exposed perithecial contours and wrinkles. Externally black, interior white. Texture soft. *Perithecia* subglobose, 0.4–0.6 mm diam.



Figure 4. *Xylaria polysporicola* (FCATAS 848, holotype) **a, b** stromata on leaves (b, FCATAS 851) **c** stromatal surface **d** section through stroma, showing perithecia **e, g** asci and ascal apical ring in Melzer's reagent **f, i** ascal apical ring in Melzer's reagent **h** asci in black India ink **j** ascospore with germ slit in 1% SDS **k, l** ascospore in water **m, n** ascospore showing a slimy sheath and non-cellular appendages in India ink (FCATAS 850) **o** Ascospore in 1% SDS. Scale bars: 1 cm (**a, b**); 0.2 mm (**c, d**); 10 μm (**e–o**).

Ostioles papillate. *Asci* with eight ascospores arranged in uniseriate manner, cylindrical, 115–185 μm total length, the spore-bearing parts 75–100 μm long × 6.5–9 μm broad, the stipes 30–90 μm long, with apical apparatus bluing in Melzer's reagent, inverted hat-shaped or urn-shaped, 2.5–4.5 μm high × 2–3.2 μm broad. *Ascospores* brown to

dark-brown, unicellular, ellipsoidal-inequilateral, with broadly rounded ends, one end slightly pinched sometimes, smooth, (11.5–)12.5–14.5(–15) × 5.5–8 μm (M = 13.2 × 6.4 μm, n=60), with straight germ slit slightly less than spore-length, a slimy sheath or non-cellular appendages visible occasionally in Indian ink.

Additional specimens examined. China. Hainan Province, Lingshui County, Diaoluoshan Natural Reserve, on fallen leaves of *Polyspora hainanensis*, 31 December 2020, Haixia Ma (FCATAS 849); 5 July 2019, Haixia Ma (FCATAS 850 & 851).

Remarks. *Xylaria polysporicola* is morphologically similar to *X. phyllocharis* Mont. However, *X. phyllocharis* has fully immersed perithecia, the fertile part with peg-like structures and smaller perithecia 0.2–0.3 mm diam (San Martín and Rogers 1989; Fournier et al. 2020). *Xylaria polysporicola* is similar to *Xylaria* sp. (80082005) from Taiwan in stromatal morphology, but the latter has slightly smaller stroma (11–14 mm total length × 1 mm diam. vs. 10–40 mm total length × 0.5–1.6 mm diam.), hard texture, slightly larger ascospores 13.5–16.5 × 5–6 μm, with narrowly rounded ends (Ju and Rogers 1999). *Xylaria phyllophila* Ces. somewhat resembles *X. polysporicola* in stromatal morphology, but the former has larger ascospores 20 × 10 μm (Cooke 1883).

Xylaria polysporicola is somewhat similar to *X. amphithele* F. San Martín & J.D. Rogers in shape and size of apical apparatus and ascospores. However, *X. amphithele* has globose to conical stromata with 3–4 to 20 naked perithecia (San Martín and Rogers 1989). In the phylogenetic tree, *X. polysporicola* formed a lineage close to *X. amphithele* and *X. ficicola*, but is distant from *X. phyllocharis*.

Discussion

We included ten *Xylaria* species on fallen leaves in the phylogenetic analyses of the present study. Except for *X. phyllocharis*, the other nine studied species formed a monophyletic clade with two wood-inhabiting species, *X. muscula* Lloyd and *X. crinalis* Hai X. Ma, Lar. N. Vassiljeva & Yu Li, in our phylogenetic tree (Figure 1). In China, only three species have been previously reported with molecular evidence: *X. ficicola* from tropical Yunnan, *X. sicula* f. *major* from tropical Taiwan and *X. betulicola* from temperate Jilin (Ma and Li 2018). Within the clade, *X. meliacearum*, associated with petioles and infructescence of *Guarea guidonia*, formed a separate branch from other *Xylaria* species on other leaves. In Hsieh et al. (2010), *X. phyllocharis* grouped with the wood-inhabiting *Xylaria* species, which did not reveal any contradictions in our tree. Three species, *X. polysporicola*, *X. amphithele* and *X. ficicola* formed a highly supported clade. Morphologically, these species have some similar features, such as ascospores with slimy sheath or non-cellular appendages, inverted hat-shaped or urn-shaped apical apparatus (San Martín and Rogers 1989; Ma et al. 2020). As *Xylaria hedyosmicola* formed a fully supported clade with *Xylaria* sp. 6, the two species should be the same, based on the ITS-TUB-RPB2 (Hsieh et al. 2010). *Xylaria lindericola*, on leaves of *Lindera robusta* formed a sister lineage to *X. sicula* f. *major* on unknown fallen leaves with high bootstrap value 100%. *Xylaria muscula*, growing on dead branches, formed a weakly supported branch with *X. lindericola* and *X. sicula* f. *major* associated

with fallen leaves in our tree. This may be because our phylogenetic analysis did not include more taxa related to *X. muscula*.

Until now, ten taxa, *X. betulicola*, *X. diminuta* F. San Martín & J.D. Rogers, *X. ficicola*, *X. foliicola* G. Huang & L. Guo, *X. hainanensis* Y.F. Zhu & L. Guo, *X. hedyosmicola*, *X. jiangsuensis* G. Huang & L. Guo, *X. lindericola*, *X. polysporicola* and *X. sicula* f. *major* have been found on fallen leaves in China (Hsieh et al. 2010; Ma et al. 2011; Zhu and Guo 2011; Huang et al. 2014, 2015; Ma and Li 2018). Amongst these species, *X. diminuta*, originally reported from Mexico, was found in Yunnan province of China in 2013 (Huang et al. 2014). *Xylaria sicula* f. *major* was first described from Sicily in 1878 and then found in Spain, Kenya, Sardinia, and Taiwan province of China (Hsieh et al. 2010; Fournier 2014). Unfortunately, except for the three species in this study, the molecular data of the other *Xylaria* species from China were not available. We anticipate that additional species of *Xylaria* on fallen leaves will be discovered as more studies are conducted.

Key to species of *Xylaria* on fallen leaves in China

1	Stromata with rounded fertile apices **2**
–	Stromata with acute sterile apices **3**
2	Stromata associated with leaves and petioles of *Ficus auriculata* (Moraceae), ascospores (16–)17.5–21(–22.7) × 6.5–8.5 µm ..... ***X. ficicola***
–	Stromata associated with leaves of *Lindera robusta* (Lauraceae), ascospores (12.5–)13.5–15.5(–18) × (7–)7.5–8.5(–9.5) µm ..... ***X. lindericola***
3	Stipes tomentose ..... ***X. hainanensis***
–	Stipes glabrous **4**
4	Fertile part subglobose ..... ***X. sicula* f. *major***
–	Fertile part not subglobose **5**
5	Stromata cylindrical **6**
–	Stromata filiform **8**
6	Ascospores (5.5–)6–8 × 3–3.5(–4) µm ..... ***X. diminuta***
–	Ascospores length > 8.5 µm 7
7	Stromata with conspicuous perithecial contours, ascospores (11.5–)12.5–14.5(–15) × 5.5–8 µm ..... ***X. polysporicola***
–	Stromata with inconspicuous perithecial contours, ascospores (8.5–)9–11 × 4–6 µm ..... ***X. foliicola***
8	Ascospores 16.5–20(–21.5) × 4–5(–6) µm ..... ***X. jiangsuensis***
–	Ascospores length < 16.5 µm **9**
9	Stromata associated with leaves of *Betula* (Betulaceae), ascospores (11.5)12–14(15) × 5–6 µm, with a straight germ slit, without appendages visible in India ink ..... ***X. betulicola***
–	Stromata associated with leaves of *Hedyosmum orientale* (Chloranthaceae), ascospores (12–)13–15(–16.7) × (6–)6.5–7.5(–8.5) µm, with straight to slightly sigmoid germ slit, with appendages visible in Indian ink..... ***X. hedyosmicola***

Acknowledgements

The authors thank Prof. Yu-Ming Ju (Institute of Plant and Microbial Biology, Academia Sinica, Taiwan, China) for suggestions on the manuscript. This study was supported by the National Natural Science Foundation of China (no. 31770023, 31972848, U1803232). We are also grateful to the Key Research and Development Program of Hainan (ZDYF2020062) and Hainan Basic and Applied Research Project for Cultivating High-Level Talents (2019RC305).

References

Ciccarone (1947) Alcune osservazioni su una forma de *Xylaria sicula* Pass. e Beltr. Nuovo Giornale Botanico Italiano 53(1946): 356–358.

Cooke MC (1883) On Xylaria and its allies. Grevillea 11: 81–94.

Cui BK, Li HJ, Ji X, Zhou JL, Song J, Si J, Yang ZL, Dai YC (2019) Species diversity, taxonomy and phylogeny of Polyporaceae (Basidiomycota) in China. Fungal Diversity 97: 137–392. https://doi.org/10.1007/s13225-019-00427-4

Dai YC (2012) Polypore diversity in China with an annotated checklist of Chinese polypores. Mycoscience 53: 49–80. https://doi.org/10.1007/s10267-011-0134-3

Dai YC, Yang ZL, Cui BK, Yu CJ, Zhou LW (2009) Species diversity and utilization of medicinal mushrooms and fungi in China. International Journal of Medicinal Mushrooms 11: 287–302. https://doi.org/10.1615/IntJMedMushr.v11.i3.80

Dennis RWG (1956) Some Xylarias of tropical America. Kew Bulletin 11: 401–444. https://doi.org/10.2307/4109126

Fournier J (2014) Update on European species of *Xylaria*, 120 pp.

Fournier J, Lechat C, Courtecuisse R (2020) The genus *Xylaria* sensu lato (Xylar-iaceae) in Guadeloupe and Martinique (French West Indies) III. Taxa with sl-ender upright stromata. Ascomycete.org 12(3): 81–164. https://doi.org/10.1128/jb.172.8.4238-4246.1990

Gao Q, Yang ZL (2016) Diversity and distribution patterns of root-associated fungi on herbaceous plants in alpine meadows of southwestern China. Mycologia 108: 281–291. https://doi.org/10.3852/14-324

Graniti (1959) Presenza di *Xylaria sicula* Pass. Et Beltr. su frutti e foglie di olivo e considerazioni sulla specie e sulle sue forme. Nuovo Giornale Botanico Italiano 66: 364–376.

Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. NucleicAcids Symposium Series 41: 95–98. https://doi.org/10.1021/bk-1999-0734.ch008

Hashemi SA, Zare R, Khodaparast SA, Elahinia SA (2015) A new *Xylaria* speciesfrom Iran. Mycologia Iranica 2: 1–10.

Hsieh HM, Ju YM, Rogers JD (2005) Molecular phylogeny of *Hypoxylon* and rel-ated genera. Mycologia 97(4): 844–865. https://doi.org/10.1080/15572536.2006.11832776

Hsieh HM, Lin CR, Fang MJ, Rogers JD, Fournier J, Lechat C, Ju YM (2010) P-hylogenetic status of *Xylaria* subgen. *Pseudoxylaria* among taxa of the subfa-mily Xylarioideae

(Xylariaceae) and phylogeny of the taxa involved in the s-ubfamily. Molecular Phylogenetics and Evolution 54: 957–969. https://doi.org/10.1016/j.ympev.2009.12.015

Huang G, Guo L, Liu N (2014) Two new species of *Xylaria* and *X. diminuta* ne-w to China. Mycotaxon 129(1): 149–152. https://doi.org/10.5248/129.149

Huang G, Wang RS, Guo L, Liu N (2015) Three new species of *Xylaria* from China. Mycotaxon 130: 299–304. https://doi.org/10.5248/130.299

Huelsenbeck JP, Ronquist F (2005) Bayesian analysis of molecular evolution using MrBayes. In: Nielsen R (Ed.) Statistical methods in molecular evolution. Springer, New York, 183–226. https://doi.org/10.1007/0-387-27733-1_7

Ju YM, Rogers JD (1999) The Xylariaceae of Taiwan (excluding *Anthostomella*). Mycotaxon 73: 343–440.

Ju YM, Hsieh HM (2007) *Xylaria* species associated with nests of *Odontotermes f-ormosanus* in Taiwan. Mycologia 99: 936–957. https://doi.org/10.1080/15572536.2007.11832525

Ju YM, Rogers JD, Hsieh HM (2018) *Xylaria* species associated with fallen fruitsand seeds. Mycologia 110(4): 726–749. https://doi.org/10.1080/00275514.2018.1469879

Kim CS, Jo GW, Kwag YN, Oh SO, Lee SG, Sung GH, Oh G, Shrestha B, Ki-m SY, Shin CH, Han SK (2016) New Records of *Xylaria* Species in Korea: *X. ripicola* sp. nov. and *X. tentaculata*. Mycobiology 44(1): 21–28. https://doi.org/10.5941/MYCO.2016.44.1.21

Laessoe T, Lodge DJ (1994) Three host-specific *Xylaria* species. Mycologia 86(3): 436–446. https://doi.org/10.1080/00275514.1994.12026431

Liu YJ, Whelen S, Hall BD (1999) Phylogenetic relationships among ascomycetes: evidence from an RNA polymerse II subunit. Molecular Biology And Evolution 16: 1799–1808. https://doi.org/10.1093/oxfordjournals.molbev.a026092

Ma HX, Li Y (2018) *Xylaria crinalis* and *X. betulicola* from China – two new s-pecies with thread-like stromata. Sydowia 70: 37–49. https://doi.org/10.12905/0380.sydowia70-2018-0037

Ma HX, Vasilyeva L, Li Y (2011) A new species of *Xylaria* from China. Mycotaxon 116: 151–155. https://doi.org/10.5248/116.151

Ma HX, Qu Z, Peng MK, Li Y (2020) Two penzigioid *Xylaria* species described from China based on morphological and molecular characters. Phytotaxa 436(1): 036–044. https://doi.org/10.11646/phytotaxa.436.1.3

Nylander JAA (2004) *MrModeltest* 2.3. Program distributed by the author. Evolutionary Biology Center, Uppsala University.

O'Donnell K, Cigelnik E (1997) Two Divergent Intragenomic rDNA ITS2 Types within a Monophyletic Lineage of the Fungus. Molecular Phylogenetics and Evolution 7(1): 103–116. https://doi.org/10.1006/mpev.1996.0376

Persoh D, Melcher M, Graf K, Fournier J, Stadler M, Rambold G (2009) Molecu-lar and morphological evidence for the delimitation of *Xylaria hypoxylon*. M-ycologia 101(2): 256–268. https://doi.org/10.3852/08-108

Petch (1915) Annals of the Royal Botanical Gardens, Peradeniya 6(1): e68.

Roensch P, Roensch S, Reiher A, Otto P (2010) Investigations on the fructicolous *Xylaria delitschii* and *Xylaria Oxyacanthae*. Boletus 32 (2): 106–122.

Rogers JD (1986) Provisional keys to *Xylaria* species in continental United States. Mycotaxon 26: 85–97.

Rogers JD, Samuels GJ (1986) Ascomycetes of New Zealand 8. *Xylaria*. New Ze-aland Journal of Botany 24(4): 615–650. https://doi.org/10.1080/0028825X.1986.10409947

Rogers DJ, Callan BE, Samuels GJ (1987) The Xylariaceae of the rain forests of North Sulawesi (Indonesia). Mycotaxon 31: 113–172.

Rogers JD, Ju YM, Lehmann J (2005) Some *Xylaria* species on termite nests. Mycologia 97(4): 914–923. http://dx.doi.org/10.1080/15572536.2006.11832783

Rogers JD, Callan BE, Rossman AY, Samuels GJ (1988) *Xylaria* (Sphaeriales, Xylariaceae) from Cerro de la Neblina, Venezuela. Mycotaxon 31: 103–153. https://doi.org/10.1007/BF00455669

Samuels GJ, Rogerson CT (1990) New Ascomycetes from the Guayana Highland. Memoirs of the New York Botanical Garden 64: 165–183.

San Martin GF, Rogers JD (1989) A preliminary account of *Xylaria* of Mexico. Mycotaxon 34: 283–373.

Stamatakis A (2006) RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. Bioinformatics 22: 2688–2690. https://doi.org/10.1093/bioinformatics/btl446

Wendt LC, Sir EB, Kuhnert E, Heitkaemper S, Lambert C, Hladki AI, Romero A-I, Luangsa-ard JJ, Srikitikulchai P, Persoh D, Stadler M (2018) Resurrection and emendation of the Hypoxylaceae, recognised from amultigene phylogeny of the Xylariales. Mycological Progress 17: 115–154. https://doi.org/10.1007/s11557-017-1311-3.

White TJ, Burns T, Lee S, Taylor J (1990) Amplification and direct sequencing o-f fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (Eds) PCR protocols, a guide to methods andapplications. Academic, San Diego, 315–322. https://doi.org/10.1016/B978-0-12-372180-8.50042-1

Yang XB (2015) The colored illustrated flora of Hainan Province. Science Press, Beijing. [in Chinese]

Zhu YF, Guo L (2011) *Xylaria hainanensis* sp. nov. (Xylariaceae) from China. M-ycosystema 30(4): 526–528. https://doi.org/10.13346/j.mycosystema.2011.04.014